



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities of the United States.

CALIFORNIA—Los Angeles.—Month of June, 1899. Estimated population, 103,000. Total number of deaths, 116, including diphtheria, 3; enteric fever, 1; whooping cough, 1; cerebro-spinal meningitis, 2, and 20 from phthisis pulmonalis.

Oakland.—Month of June, 1899. Estimated population, 75,000. Total number of deaths, 60, including enteric fever, 3; cerebro-spinal meningitis, 1; smallpox, 1, and 4 from phthisis pulmonalis.

Sacramento.—Month of June, 1899. Estimated population, 30,000. Total number of deaths 32, including whooping cough 1, and 6 from phthisis pulmonalis.

CONNECTICUT.—Reports to the State board of health for the month of June, 1899, from 165 towns, having an aggregate population of 910,294, show a total of 1,048 deaths, including diphtheria, 7; enteric fever, 2; measles, 3; scarlet fever, 5; whooping cough, 17; la grippe, 4; cerebro-spinal meningitis, 12, and 106 from phthisis pulmonalis.

Bridgeport.—Month of June, 1899. Estimated population, 65,000. Total number of deaths, 99, including scarlet fever, 1, and 5 from whooping cough.

New Haven.—Month of June, 1899. Estimated population, 74,000. Total number of deaths, 120, including whooping cough, 1, and 8 from phthisis pulmonalis.

IOWA—Boone.—Month of May, 1899. Census population, 6,520. Total number of deaths, 6.

Burlington.—Estimated population, 30,000. Total number of deaths, 20, including 5 from phthisis pulmonalis.

Carroll.—Estimated population, 3,000. Total number of deaths, 2.

Cedar Falls.—Estimated population, 6,000. Total number of deaths, 4.

Cedar Rapids.—Estimated population, 27,000. Total number of deaths, 19, including la grippe, 1, and 1 from phthisis pulmonalis.

Cherokee.—Census population, 3,441. Total number of deaths, 1.

Davenport.—Estimated population, 40,000. Total number of deaths, 48, including enteric fever, 1, and 4 from phthisis pulmonalis.

Des Moines.—Estimated population, 72,000. Total number of deaths, 70, including diphtheria, 1; enteric fever, 1, and 5 from phthisis pulmonalis.

Eldon.—Estimated population, 2,300. Total number of deaths, 1.

Emmetsburg.—Estimated population, 2,400. Total number of deaths, 1.

Fairfield.—Census population, ———. Total number of deaths, 2.

Keokuk.—Estimated population, 19,305. Total number of deaths, 23, including la grippe, 2, and 2 from phthisis pulmonalis.

Le Mars.—Estimated population, 5,000. Total number of deaths, 8, including diphtheria, 1, and 1 from phthisis pulmonalis.

Marengo.—Estimated population, 2,500. Total number of deaths, 2, including 2 from measles.

Ottumwa.—Estimated population, 20,000. Total number of deaths, 26, including enteric fever, 1; measles, 2, and 4 from phthisis pulmonalis.

Sibley.—Estimated population, 1,500. Total number of deaths, 1.

The Health Bulletin says:

The following outbreaks of infectious disease have been reported for the month of May:

Cerebro-spinal meningitis.—Linn Grove; Ottumwa; Davenport.

Diphtheria.—Dows; East Orange Township, Sioux County; Liberty Township, Hamilton County; Riceville; Alton; Harrison Township, Harrison County; Dunlap; Linn Township, Linn County; Des Moines; Dubuque; Le Mars.

Measles.—Sibley; Rockford; Dubuque; Marengo.

Scarlet fever.—Alton; Correctionville; Lynnvile; Kamrar; Harcourt; La Porte City; Lost Grove Township, Webster County; Liberty Township, Hamilton County; Wapello; Liscomb.

Smallpox.—Cresco; Rome; Orleans Township, Winneshiek County; Le Claire.

There are at present, as reported since May, cases in Pleasant Valley, Scott County, 1 case at Iowa City and cases in three families in Paris Township, Howard County. The cases in Pleasant Valley Township and Iowa City resulted from exposure at Le Claire, and those in Paris Township from exposure in Cresco.

LATER: Dr. Shrader reports, June 26, another case of smallpox at Iowa City. The city has been enjoined from using a pesthouse erected by it in a township outside the city limits.

Typhoid fever.—Davenport; Des Moines; Ottumwa.

Whooping cough.—Slater; Keokuk.

LOUISIANA—*New Orleans*.—Month of June, 1899. Estimated population, 300,000—white, 210,000; colored, 90,000. Total number of deaths, 738—white, 458; colored, 280, including enteric fever, 21; measles, 1; cerebro-spinal meningitis, 10; smallpox, 1, and 77 from phthisis pulmonalis.

MARYLAND—*Baltimore*.—Month of May, 1899. Estimated population, 541,000—white, 463,000; colored, 78,000. Total number of deaths, 752—white, 553; colored, 197, including diphtheria, 9; enteric fever, 3; scarlet fever, 3; influenza, 6; cerebro-spinal meningitis, 16, and 85 from phthisis pulmonalis.

MASSACHUSETTS—*Brockton*.—Month of June, 1899. Estimated population, 37,278. Total number of deaths, 36, including 6 from phthisis pulmonalis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended July 8, 1899, from 97 observers, indicate that measles, dysentery, inflammation of kidney, and inflammation of bowels increased and cerebro-spinal meningitis and intermittent fever decreased in area of prevalence. Phthisis pulmonalis was reported present at 182, measles at 48, enteric fever at 35, scarlet fever at 31, whooping cough at 18, diphtheria at 13, and cerebro-spinal meningitis at 5 places.

MINNESOTA—*St. Paul*.—Month of June, 1899. Estimated population, 200,000. Total number of deaths, 120, including diphtheria, 4; enteric fever, 1; whooping cough, 2, and 16 from phthisis pulmonalis.

Winona.—Month of June, 1899. Estimated population, 24,000. Total number of deaths, 16, including cerebro-spinal meningitis, 1, and 1 from phthisis pulmonalis.

MISSOURI—*St. Louis*.—Month of May, 1899. Estimated population, 623,000—white, 587,000; colored, 36,000. Total number of deaths, 775—white, 675; colored, 100, including diphtheria, 17; enteric fever, 2; measles, 2; whooping cough, 7, and 91 from phthisis pulmonalis.

NEW HAMPSHIRE—*Concord*.—Month of June, 1899. Estimated population, 18,000. Total number of deaths, 30, including 2 from phthisis pulmonalis.

Manchester.—Month of June, 1899. Estimated population, 60,000. Total number of deaths, 76, including scarlet fever, 1; whooping cough, 1, and 5 from phthisis pulmonalis.

NEW YORK—*Buffalo*.—Month of June, 1899. Estimated population, 370,000. Total number of deaths, 305, including diphtheria, 7; enteric fever, 5; measles, 3; scarlet fever, 1; whooping cough, 2; cerebro-spinal meningitis, 3, and 33 from phthisis pulmonalis.

Yonkers.—Month of June, 1899. Estimated population, 43,000. Total number of deaths, 57, including diphtheria, 1; scarlet fever, 1, and 8 from phthisis pulmonalis.

OHIO—*Cleveland*.—Month of June, 1899. Estimated population, 395,000. Total number of deaths, 394, including diphtheria, 3; enteric fever, 10; scarlet fever, 2; cerebro-spinal meningitis, 5; smallpox, 1; whooping cough, 4, and 16 from phthisis pulmonalis.

PENNSYLVANIA—*Meadville*.—Month of June, 1899. Estimated population, 12,000. Total number of deaths, 7. No deaths from contagious diseases.

Williamsport.—Month of June, 1899. Estimated population, 35,000. Total number of deaths, 20, including scarlet fever, 1, and 4 from phthisis pulmonalis.

TEXAS—*Galveston*.—Month of March, 1899. Estimated population, 60,000. Total number of deaths, 74, including diphtheria 2; cerebro-spinal meningitis 3, and 2 from phthisis pulmonalis.

Month of April. Total number of deaths, 62, including diphtheria, 1; cerebro spinal meningitis, 5, and 3 from phthisis pulmonalis.

Month of May. Total number of deaths, 62, including measles, 2; cerebro spinal meningitis, 2, and 2 from phthisis pulmonalis.

San Antonio.—Month of June, 1899. Estimated population, 63,000. Total number of deaths, 166, including enteric fever, 9; measles, 5, and 19 from phthisis pulmonalis, mostly nonresidents.

WISCONSIN—*Superior*.—Month of June, 1899. Estimated population, 30,000. Total number of deaths, 37, including diphtheria, 2, and 2 from phthisis pulmonalis.